



## FTCD gene

formiminoyltransferase cyclodeaminase

### Normal Function

The *FTCD* gene provides instructions for making an enzyme called formiminotransferase cyclodeaminase. This enzyme is found mainly in the liver, with smaller amounts in the kidneys. In males it is also found in the testes.

Formiminotransferase cyclodeaminase is called a bifunctional enzyme because it performs two functions. Specifically, it is involved in the last two steps in the breakdown (metabolism) of the amino acid histidine, a building block of most proteins. It also plays a role in producing one of several forms of the vitamin folate, which has many important functions in the body.

One of the enzyme's functions, called formiminotransferase, breaks down a molecule called N-formiminoglutamate in the process of histidine metabolism. Part of the N-formiminoglutamate molecule, called the formimino group, is incorporated into another molecule called formiminotetrahydrofolate. The amino acid glutamate is also produced in this reaction.

The second function of the enzyme, called cyclodeaminase, breaks down formiminotetrahydrofolate to a molecule called 5,10-methenyltetrahydrofolate and ammonia. 5,10-methenyltetrahydrofolate is one of several tetrahydrofolate molecules involved in the production (synthesis) of many important molecules in the body, such as purines and pyrimidines (the building blocks of DNA and its chemical cousin, RNA) and amino acids. The tetrahydrofolates carry small molecules called one-carbon units that are needed for the synthesis of these larger molecules.

### Health Conditions Related to Genetic Changes

#### glutamate formiminotransferase deficiency


At least three *FTCD* gene mutations that cause glutamate formiminotransferase deficiency have been identified. Two of these mutations change an amino acid in the formiminotransferase cyclodeaminase enzyme sequence. These mutations reduce one of the enzyme's two activities, formiminotransferase. Another mutation results in the production of a formiminotransferase cyclodeaminase enzyme that is shortened so that the part of the enzyme that takes part in the cyclodeaminase reaction is missing. Disruption of these reactions results in the excretion of a molecule called formiminoglutamate (FIGLU) in the urine of affected individuals and, in some severe cases of glutamate formiminotransferase deficiency, a buildup of certain B vitamins

called folates in the blood. It is unclear how the mutations result in the specific health problems associated with glutamate formiminotransferase deficiency.

### Chromosomal Location

Cytogenetic Location: 21q22.3, which is the long (q) arm of chromosome 21 at position 22.3

Molecular Location: base pairs 46,135,981 to 46,156,481 on chromosome 21 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

### Other Names for This Gene

- FTCD\_HUMAN
- LCHC1

### Additional Information & Resources

#### Educational Resources

- Biochemistry (fifth edition, 2002): Tetrahydrofolate Carries Activated One-Carbon Units at Several Oxidation Levels  
<https://www.ncbi.nlm.nih.gov/books/NBK22459/#A3357>

#### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28FTCD%5BTIAB%5D%29%29+OR+%28formiminotransferase+cyclodeaminase%5BTIAB%5D%29%29+OR+%28formimidoyltransferase+cyclodeaminase%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

#### OMIM

- FORMIMINOTRANSFERASE CYCLODEAMINASE  
<http://omim.org/entry/606806>

## Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology  
[http://atlasgeneticsoncology.org/Genes/GC\\_FTCD.html](http://atlasgeneticsoncology.org/Genes/GC_FTCD.html)
- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=FTCD%5Bgene%5D>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=3974](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=3974)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/10841>
- UniProt  
<http://www.uniprot.org/uniprot/O95954>

## **Sources for This Summary**

- OMIM: FORMIMINOTRANSFERASE CYCLODEAMINASE  
<http://omim.org/entry/606806>
- Hilton JF, Christensen KE, Watkins D, Raby BA, Renaud Y, de la Luna S, Estivill X, MacKenzie RE, Hudson TJ, Rosenblatt DS. The molecular basis of glutamate formiminotransferase deficiency. *Hum Mutat.* 2003 Jul;22(1):67-73. Erratum in: *Hum Mutat.* 2003 Nov;22(5):416.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12815595>
- Mao Y, Vyas NK, Vyas MN, Chen DH, Ludtke SJ, Chiu W, Quijcho FA. Structure of the bifunctional and Golgi-associated formiminotransferase cyclodeaminase octamer. *EMBO J.* 2004 Aug 4;23(15):2963-71. Epub 2004 Jul 22.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15272307>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC514939/>
- Solans A, Estivill X, de la Luna S. Cloning and characterization of human FTCD on 21q22.3, a candidate gene for glutamate formiminotransferase deficiency. *Cytogenet Cell Genet.* 2000;88(1-2):43-9.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/10773664>

---

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/gene/FTCD>

Reviewed: August 2009  
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services